


EXHIBIT D



Entrez

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

B

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Comput Chem. 2001 Jul; 25(4): 411-22.

Related Articles, L

ELSEVIER
FULL-TEXT ARTICLE**Amino acid residue environments and predictions of residue ty****Kim W, Wilbur WJ.**

National Center for Biotechnology Information, National Library of Medicine
National Institutes of Health, Bethesda, MD 20894, USA.
wonkim@ncbi.nlm.nih.gov

The determination of a protein's structure from the knowledge of its linear chain is one of the important problems that remains as a bottleneck in interpreting the rapidly increasing repository of genetic sequence data. One approach to this problem that has shown promise and given a measure of success is threading. In this approach contact energies between different amino acids are first determined by statistical methods applied to known structures. These contact energies are then applied to a sequence whose structure is to be determined by threading it through various known structures and determining the total threading energy for each candidate structure. That structure that yields the lowest total energy is then considered the leading candidate among all the structures tested. Additional information is often needed in order to support the results of threading studies, as it is well known in the field that contact potentials used are not sufficiently sensitive to allow definitive conclusions. Here, we investigate the hypothesis that the environment of an amino acid residue realized as all those residues not local to it on the chain but sufficiently close spatially can supply information predictive of the type of residue that is not adequately reflected in the individual contact energies. We present evidence that confirms this hypothesis and suggests a high order cooperativity between the residues that surround a given residue and how they interact with it. We suggest a possible application to threading.

PMID: 11459355 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)